



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

York. While it would be ill advised to propose a revival of *C. scabrior* on the evidence at command, attention may well be drawn to the very probable distinctness of the two plants.

### Two Californian Saxifrages.

#### SAXIFRAGA FALLAX.

Perennial, fibrous-rooted, propagating by abundant subterranean short rhizomatous offsets; stem scapiform, 8-12 inches high, loosely cymose-paniculate at summit, the whole minutely and sparsely glandular-hispidulous, but with no villous hairs; leaves thin, ovate or oval, subcordate, or truncate at base, saliently toothed or only repand; calyx very deeply cleft, the segments reflexed; petals white, obovate-oblong, obtuse, emarginate; filaments very short; anthers green; mature carpels divergent almost from the base.

A northeast Californian and subalpine ally of the widely distributed *S. Californica*, differing from that in its short subterranean offsets, its emarginate petals and almost completely divergent carpels. I have it only from Lassen's Peak, where it was collected by Mrs. Austin long since, and from above Donner Lake, where I obtained it last summer.

SAXIFRAGA APRICA. *S. umbellulata* Greene, Eryth. 1: 222. The name first assigned to this small, but common and well marked species of sunny exposures near the crest of the Sierra, must yield to the much earlier *S. umbellulata* of Hook. f. & Thoms.

EDW. L. GREENE.

### Proceedings of the Club.

ANNUAL MEETING, TUESDAY EVENING, JANUARY 14TH, 1896.

The President occupied the chair and there were 39 persons present.

The following were elected active members: Miss Mary Edgar, Mr. S. Sidney Smith, Mrs. S. Sidney Smith, Miss Mary Appleton, Miss Mary Farley, Miss Laura H. Knupfer, Mr. C. L. Allen, Mr. B. A. Gilbert, Mr. Per Axel Rydberg, Mr. Oliver A. Farwell.

The Finance Committee reported through the Secretary that no business had been before them for transaction during the year.